

## § 77.34

## 9 CFR Ch. I (1–1–01 Edition)

the test is administered or by a veterinarian employed by USDA.

(1) A designated accredited veterinarian may conduct the SCT test, except as provided in § 77.34(a)(2) and § 77.39(e) and (f).

(2) Any accredited veterinarian may conduct the BTB test.

(b) *Approved diagnostic laboratories.*

(1) With one exception, histopathology and culture results for all tuberculosis diagnoses will be accepted only from the National Veterinary Services Laboratories (NVSL) in Ames, IA. The exception is that results will be accepted from a laboratory of the Food Safety and Inspection Service, USDA, for tissue examination of regular-kill slaughter animals in those cases where no submission is made to NVSL.

(2) The following laboratory is approved to perform the BTB test: Texas Veterinary Medical Center laboratory at Texas A&M University in College Station, TX.

(c) *Identification.* Any captive cervid tested with an official tuberculosis test must bear official identification in the form of an official eartag, or another identification device or method approved by the Administrator as unique and traceable, at the time of the official tuberculosis test. Use of any identification device or method other than an official eartag must first be approved by the Administrator as unique and traceable. Written requests for approval must be sent to National Animal Health Programs, VS, APHIS, 4700 River Road Unit 43, Riverdale, MD 20737–1231.

(d) *Reporting of tests.*

(1) *SCT and CCT tests.* For the SCT and CCT tests, the testing veterinarian must submit a report to cooperating State and Federal animal health officials of the State in which the captive cervid is tested. The report must include the following information for all SCT and CCT tests administered: The number of the individual eartag or other identification approved by the Administrator; the age, sex, and breed of each captive cervid tested; a record of all responses; the size of each response for the CCT test; and the test interpretation.

(2) *BTB test.* Copies of the BTB test results must be submitted by the testing laboratory to the person, firm, or corporation responsible for the management of the herd, cooperating State and Federal animal health officials of the State in which the captive cervid is tested, and the testing veterinarian. The report must include the following information for all BTB tests administered: The number of the individual eartag or other identification approved by the Administrator; the age, sex, and breed of each captive cervid tested; the test interpretation, and a summary of supporting data. Full supporting data must be submitted by the testing laboratory on a case-by-case basis at the request of cooperating State and Federal animal health officials.

(e) *Test interpretation.*

(1) Interpretation of an SCT test will be based upon the judgment of the testing veterinarian after observation and palpation of the injection site, in accordance with the classification requirements described in § 77.34(a).

(2) Interpretation of a CCT test will be in accordance with the classification requirements described in § 77.34(b).

(3) Interpretation of a BTB test will be in accordance with the patented standards for the BTB test<sup>2</sup> and the classification requirements described in § 77.34(c).

(f) *Captive cervids eligible for testing.* Except as provided in § 77.35(a)(1) and § 77.36(a)(1), testing of herds for individual herd classification must include all captive cervids 1 year of age or over and any captive cervids other than natural additions (captive cervids born into the herd) under 1 year of age.

### § 77.34 Official tuberculosis tests.

(a) *Single cervical tuberculin (SCT) test.*

(1) The SCT test is the primary test to be used in individual captive cervids and in herds of unknown tuberculous

<sup>2</sup>The patented standards for the BTB test may be obtained from the Texas Veterinary Medical Center, College of Veterinary Medicine, Texas A&M University, College Station, TX, or from the Deer Research Laboratory, Department of Microbiology, University of Otago, P.O. Box 56, Dunedin, New Zealand.

status. Each captive cervid that responds to the SCT test must be classified as a suspect until it is retested with either the CCT test or the BTB test and is either found negative for tuberculosis or is classified as a reactor, unless, with the exception of a designated accredited veterinarian, the testing veterinarian determines that the captive cervid should be classified as a reactor based on its response to the SCT test. A designated accredited veterinarian must classify a responding captive cervid as a suspect, unless the DTE determines, based on epidemiological evidence, that the captive cervid should be classified as a reactor.

(2) The SCT test is the primary test to be used in affected herds and in herds that have received captive cervids from an affected herd. When used with affected herds or in herds that have received captive cervids from an affected herd, the SCT test may only be administered by a veterinarian employed by the State in which the test is administered or employed by USDA. In affected herds or herds that have received captive cervids from an affected herd, each captive cervid that responds to the SCT test must be classified as a reactor, unless the DTE determines that the captive cervid should be classified as a suspect because of possible exposure to a tuberculous animal.

(b) *Comparative cervical tuberculin (CCT) test.*

(1) The CCT test is a supplemental test that may only be used for retesting captive cervids classified as suspects. The CCT test may be used in affected herds only after the herd has tested negative to at least two whole herd SCT tests and only with the prior written consent of the DTE. The CCT test may not be used as a primary test for herds of unknown tuberculous status.

(2) A captive cervid tested with the CCT test must be classified as negative if it has a response to the bovine PPD tuberculin that is less than 1 mm.

(3) Unless the testing veterinarian determines that the captive cervid should be classified as a reactor because of possible exposure to a tuberculous animal, a captive cervid tested

with the CCT test must be classified as a suspect if:

(i) It has a response to the bovine PPD tuberculin that is greater than 2 mm and that is equal to the response to the avian PPD tuberculin; or

(ii) It has a response to the bovine PPD tuberculin that is equal to or greater than 1mm and equal to or less than 2mm and that is equal to or greater than the response to the avian PPD tuberculin.

(4) A captive cervid tested with the CCT test must be classified as a reactor if:

(i) It has a response to the bovine PPD tuberculin that is greater than 2 mm and that is at least 0.5 mm greater than the response to the avian PPD tuberculin; or

(ii) It has been classified as a suspect on two successive CCT tests.

(iii) Any exceptions to reactor classification under the conditions in paragraph (b)(4)(i) and (b)(4)(ii) of this section must be justified by the testing veterinarian in writing and have the concurrence of the DTE.

(c) *Blood tuberculosis (BTB) test.*

(1) The BTB test is a supplemental test that may be used in place of the CCT test for retesting captive cervids classified as suspects.

(2) Except as provided in § 77.39(e), any captive cervid classified by the testing laboratory as "equivocal" will be classified as a suspect.

(3) Any captive cervid classified by the testing laboratory as "*M. bovis* positive" will be classified as a reactor.

(4) Any captive cervid classified by the testing laboratory as "avian" or "negative" will be considered negative for tuberculosis.

(5) The owner of the captive cervid tested is responsible for the cost of the BTB test.

**§ 77.35 Interstate movement from accredited herds.**

(a) *Qualifications.* To be recognized as an accredited herd:

(1) All captive cervids in the herd eligible for testing in accordance with § 77.33(f) must have tested negative to at least three consecutive official tuberculosis tests, conducted at 9-15 month intervals. However, captive cervids under 1 year of age that are not