

§ 520.1840

equivalent to 17, 34, or 230 grams of piperazine per pound or 100 milliliters.

(b) *Sponsor*. See 015565 in § 510.600(c) of this chapter.

(c) *Related tolerances*. See § 556.513 of this chapter.

(d) *Conditions of use*—(1) *Chickens*—(i) *Amount*. 50 milligrams per bird under 6 weeks, 100 milligrams per bird over 6 weeks.

(ii) *Indications for use*. For removal of large roundworm (*Ascaridia* spp.).

(iii) *Limitations*. For use in drinking water or feed. Use as sole source of drinking water. Prepare fresh solution daily. Use as 1-day single treatment. Withdraw 14 days prior to slaughter. Do not use for chickens producing eggs for human consumption. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(2) *Turkeys*—(i) *Amount*. 100 milligrams per bird up to 12 weeks and 200 milligrams per bird over 12 weeks.

(ii) *Indications for use*. For removal of large roundworm (*Ascaridia* spp.).

(iii) *Limitations*. For use in drinking water or feed. Use as sole source of drinking water. Prepare fresh solution daily. Use as 1-day single treatment. Withdraw 14 days prior to slaughter. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(3) *Swine*—(i) *Amount*. 50 milligrams per pound of body weight.

(ii) *Indications for use*. For removal of large roundworm (*Ascaris suum*) and nodular worms (*Oesophagostomum* spp.).

(iii) *Limitations*. For use in drinking water or feed. Use as sole source of drinking water. Prepare fresh solution daily. Use as 1-day single treatment. Withdraw 21 days prior to slaughter. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[64 FR 23018, Apr. 29, 1999]

§ 520.1840 Poloxalene.

(a) *Specifications*. Polyoxypropylene-polyoxyethylene glycol nonionic block polymer.

(b) *Sponsors*. See sponsors in § 510.600(c) of this chapter for use as in paragraph (d) of this section.

(1) No. 000069 for use as in paragraphs (d)(1) and (d)(3) of this section.

21 CFR Ch. I (4–1–03 Edition)

(2) No. 017800 for use as in paragraph (d)(4) of this section.

(3) No. 036904 for use as in paragraph (d)(2) of this section.

(4) No. 066104 for use as in paragraph (d)(3) of this section.

(c) [Reserved]

(d) *Conditions of use*. (1) For treatment of legume (alfalfa, clover) bloat in cattle. Administer as a drench at the rate of 25 grams for animals up to 500 pounds and 50 grams for animals over 500 pounds of body weight.

(2) For control of legume (alfalfa, clover) bloat in cattle. Administer, in molasses block containing 6.6 percent poloxalene, at the rate of 0.8 oz. of block (1.5 grams poloxalene) per 100 lbs. of body weight per day.

(3) For prevention of legume (alfalfa, clover) and wheat pasture bloat in cattle. A 53-percent poloxalene top dressing on individual rations of ground feed. Dosage is 1 gram of poloxalene per 100 pounds of body weight daily. If bloating conditions are severe, the dose is doubled. Treatment should be started 2 to 3 days before exposure to bloat-producing conditions. Repeat use of the drug if animals are exposed to bloat-producing conditions for more than 12 hours after the last treatment. Do not exceed the double dose in any 24-hour period.

(4) For control of legume (alfalfa, clover) and wheat pasture bloat in cattle. Administer in molasses block containing 6.6 percent poloxalene, at the rate of 0.8 ounce of block (1.5 grams of poloxalene) per 100 pounds of body weight per day. Provide access to blocks at least 7 days before exposure to bloat-producing conditions.

[40 FR 13838, Mar. 27, 1975, as amended at 40 FR 39857, Aug. 29, 1975; 42 FR 41854, Aug. 19, 1977; 50 FR 5385, Feb. 8, 1985; 54 FR 33501, Aug. 15, 1989; 56 FR 50653, Oct. 8, 1991; 58 FR 26523, May 4, 1993; 60 FR 55659, Nov. 2, 1995; 66 FR 47963, Sept. 17, 2001]

§ 520.1846 Polyoxyethylene (23) lauryl ether blocks.

(a) *Specifications*. Each molasses-based block contains 2.2 percent polyoxyethylene (23) lauryl ether.

(b) *Sponsor*. See No. 050112 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount*. 2 grams of polyoxyethylene (23) lauryl

ether per 100 kilograms of body weight per day (1 pound of block per 500 kilogram (1,100 pound) animal per day).

(2) *Indications for use.* For reduction of the incidence of bloat (alfalfa and clover) in pastured cattle.

(3) *Limitations.* Administer free-choice to beef cattle and nonlactating dairy cattle only. Initially, provide one block per five head of cattle. Start treatment 10 to 14 days before exposure to bloat-producing pastures. Do not allow cattle access to other sources of salt while being fed this product. Do not feed this product to animals without adequate forage/roughage consumption.

[50 FR 48189, Nov. 22, 1985, as amended at 56 FR 9841, Mar. 8, 1991]

§ 520.1855 Ponazuril.

(a) *Specifications.* Each gram of paste contains 150 milligrams (mg) ponazuril.

(b) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use in horses—(1) Amount.* 5 mg per kilogram body weight, daily for 28 days.

(2) *Indications for use.* For the treatment of equine protozoal myeloencephalitis caused by *Sarcocystis neurona*.

(3) *Limitations.* Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[66 FR 43774, Aug. 21, 2001]

§ 520.1870 Praziquantel tablets.

(a) *Specifications.* Each dog tablet contains 34 milligrams (mg) of praziquantel; each cat tablet contains 11.5 or 23 mg of praziquantel.

(b) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Dogs—(i) Indications for use.* For removal of canine cestodes *Dipylidium caninum* and *Taenia pisiformis*. If labeled for use by or on the order of a licensed veterinarian, for removal of the canine cestode *Echinococcus granulosus*, and for removal and control of the canine cestode *Echinococcus multilocularis*.

(ii) *Dosage.* Dogs 5 pounds and under, ½ tablet (17 mg); 6 to 10 pounds, 1 tablet (34 mg); 11 to 15 pounds, 1½ tablets (51 mg); 16 to 30 pounds, 2 tablets (68

mg); 31 to 45 pounds, 3 tablets (102 mg); 46 to 60 pounds, 4 tablets (136 mg); over 60 pounds, 5 tablets maximum (170 mg).

(iii) *Limitations.* Administer directly by mouth or crumbled and in feed. Not intended for use in puppies less than 4 weeks of age. For over-the-counter (OTC) use: Consult your veterinarian before administering tablets to weak or debilitated animals, and for assistance in the diagnosis, treatment, and control of parasitism. For prescription use: Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats—(i) Indications for use.* For removal of feline cestodes *Dipylidium caninum* and *Taenia taeniaeformis*.

(ii) *Dosage.* Cats 4 pounds and under, 11.5 mg; 5 to 11 pounds, 23 mg; over 11 pounds, 34.5 mg.

(iii) *Limitations.* Administer directly by mouth or crumbled and in feed. Not intended for use in kittens less than 6 weeks of age. For OTC use: Consult your veterinarian before administering tablets to weak or debilitated animals, and for assistance in the diagnosis, treatment, and control of parasitism.

[46 FR 60570, Dec. 11, 1981, as amended at 47 FR 26377, June 18, 1982; 55 FR 2234, Jan. 23, 1990; 58 FR 7864, Feb. 10, 1993; 58 FR 42853, Aug. 12, 1993]

§ 520.1871 Praziquantel/pyrantel pamoate tablets.

(a) *Specifications.* Each cat tablet contains 18.2 milligrams (mg) praziquantel with 72.6 mg pyrantel (as pyrantel pamoate).

(b) *Sponsor.* See 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use—(1) Cats—(i) Dosage.* 1.5 to 1.9 pounds, ¼ tablet; 2 to 3 pounds, ½ tablet; 4 to 8 pounds, 1 tablet; 9 to 12 pounds, 1 ½ tablets; 13 to 16 pounds, 2 tablets.

(ii) *Indications for use.* For removal of tapeworms (*Dipylidium caninum* and *Taenia taeniaeformis*), hookworms (*Ancylostoma tubaeforme*), and large roundworms (*Toxocara cati*) in cats and kittens.

(iii) *Limitations.* Not for use in kittens less than 1 month of age or weighing less than 1.5 pounds. May be given directly by mouth or in a small amount of food. Do not withhold food