

(2) It is administered by stomach tube or as a drench at a dose of 0.9 gram of cambendazole per 100 pounds of body weight (20 milligrams per kilogram).

(3) For animals maintained on premises where reinfection is likely to occur, re-treatments may be necessary. For most effective results, re-treat in 6 to 8 weeks.

(4) Not for use in horses intended for food.

(5) Caution: Do not administer to pregnant mares during first 3 months of pregnancy.

(6) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13833, Mar. 27, 1975. Redesignated at 41 FR 1276, Jan. 7, 1976, and amended at 42 FR 3838, Jan. 21, 1977; 62 FR 63270, Nov. 28, 1997]

§ 520.300b Cambendazole pellets.

(a) *Specifications.* The drug is in feed pellets containing 5.3 percent cambendazole.

(b) *Sponsor.* No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in horses for the control of large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*); small strongyles (*Trichonema*, *Poteriostomum*, *Cylicobrachytus*, *Craterostomum*, *Oesophagodontus*); roundworms (*Parascaris*); pinworms (*Oxyuris*); and threadworms (*Strongyloides*).

(2) Administer 20 milligrams cambendazole per kilogram body weight (6 ounces per 1,000 pounds) by mixing with normal grain ration given at one feeding. Doses for individual horses should be mixed and fed separately to assure that each horse will consume the correct amount.

(3) For animals maintained on premises where reinfection is likely to occur, re-treatments may be necessary. For most effective results, re-treat in 6 to 8 weeks.

(4) Not for use in horses intended for food.

(5) Caution: Do not administer to pregnant mares during first 3 months of pregnancy.

(6) Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[41 FR 1276, Jan. 7, 1976, as amended at 42 FR 3838, Jan. 21, 1977; 62 FR 63270, Nov. 28, 1997]

§ 520.300c Cambendazole paste.

(a) *Specifications.* The drug is a paste containing 45 percent cambendazole.

(b) *Sponsor.* No. 050604 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in horses for the control of large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*); small strongyles (*Trichonema*, *Poteriostomum*, *Cylicobrachytus*, *Craterostomum*, *Oesophagodontus*); roundworms (*Parascaris*); pinworms (*Oxyuris*); and threadworms (*Strongyloides*).

(2) Administer 20 milligrams cambendazole per kilogram body weight (5 grams per 550 pounds (250 kilograms)) by depositing the paste on the back of the tongue using a dosing gun.

(3) For animals maintained on premises where reinfection is likely to occur, re-treatments may be necessary. For most effective results, re-treat in 6 to 8 weeks.

(4) Not for use in horses intended for food.

(5) Caution: Do not administer to pregnant mares during first 3 months of pregnancy.

(6) Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[41 FR 1276, Jan. 7, 1976, as amended at 42 FR 3838, Jan. 21, 1977; 62 FR 63270, Nov. 28, 1997]

§ 520.309 Carprofen.

(a) *Specifications.* (1) Each caplet contains 25, 75, or 100 milligrams (mg) carprofen.

(2) Each chewable tablet contains 25, 75, or 100 mg carprofen.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use in dogs*—(1) *Amount.* 2 mg per pound (1/16) of body weight once daily or 1 mg/lb twice daily. For the control of postoperative pain, administer approximately 2 hours before the procedure.

§ 520.310

(2) *Indications for use.* For the relief of pain and inflammation associated with osteoarthritis, and for the control of postoperative pain associated with soft tissue and orthopedic surgery.

(3) *Limitations.* Federal Law restricts this drug to use by or on the order of a licensed veterinarian.

[61 FR 66581, Dec. 18, 1996, as amended at 64 FR 32181, June 16, 1999; 66 FR 63165, Dec. 5, 2001; 67 FR 6866, Feb. 14, 2002; 67 FR 65038, Oct. 23, 2002; 67 FR 65697, Oct. 28, 2002]

§ 520.310 Caramiphen ethanedisulfonate and ammonium chloride tablets.

(a) *Specifications.* Each tablet contains 10 milligrams of 5stcaramiphen ethanedisulfonate and 80 milligrams of ammonium chloride.¹

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs*—(1) *Amount.* One tablet per 15 to 30 pounds of body weight every 4 to 6 hours.¹

(2) *Indications for use.* For relief of cough.¹

[43 FR 55385, Nov. 28, 1978]

§ 520.312 Carnidazole tablets.

(a) *Specifications.* Each tablet contains 10 milligrams of carnidazole.

(b) *Sponsor.* See 053923 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* Adult pigeons: 1 tablet (10 milligrams); newly weaned pigeons: ½ tablet (5 milligrams).

(2) *Indications for use.* For treating trichomoniasis (canker) in ornamental and homing pigeons.

(3) *Limitations.* Not for use in pigeons intended for human food. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism or when severely ill birds do not respond to treatment.

[54 FR 32336, Aug. 7, 1989]

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

21 CFR Ch. I (4–1–03 Edition)

§ 520.314 Cefadroxil tablets.

(a) *Specifications.* 50-, 100-, and 200-milligram tablets for dogs and cats; 1 gram tablet for dogs.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) For use in dogs as follows:

(i) *Indications for use.* For the treatment of skin and soft tissue infections including cellulitis, pyoderma, dermatitis, wound infections, and abscesses due to susceptible strains of *Staphylococcus aureus*. For the treatment of genitourinary tract infections (cystitis) due to susceptible strains of *Escherichia coli*, *Proteus mirabilis*, and *Staphylococcus aureus*.

(ii) *Amount.* Ten milligrams per pound of body weight twice daily.

(iii) *Limitations.* The drug is administered orally. For skin and soft tissue infections, treatment should be continued for a minimum of 3 days. For genitourinary tract infections, treatment should be continued for a minimum of 7 days. Continue treatment at least 48 hours after the dog has become afebrile or asymptomatic. If no response is seen after 3 days of treatment, therapy should be discontinued and the case reevaluated. Do not treat for more than 30 days. Safety for use in pregnant bitches and stud dogs has not been determined. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) For use in cats as follows:

(i) *Indications for use.* For the treatment of skin and soft tissue infections including abscesses, wound infections, cellulitis, and dermatitis caused by susceptible strains of *Pasteurella multocida*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, and *Streptococcus* spp.

(ii) *Amount.* Ten milligrams per pound of body weight once daily.

(iii) *Limitations.* The drug is administered orally. Continue treatment at least 48 hours after the cat has become afebrile or asymptomatic. If no response is seen after 3 days of treatment, therapy should be discontinued and the case reevaluated. Do not treat for more than 21 days. Safety for use in pregnant cats and breeding male cats has not been determined. Federal law