

(b) *Sponsor*. See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) The drug is used in dogs to control bacterial diarrhea caused by organisms susceptible to neomycin and to reduce smooth muscle contractions.¹

(2) It is administered at a dosage level of one to two tablets per 10 pounds of body weight twice daily for 3 days.¹

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13838, Mar. 27, 1975, as amended at 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61624, Nov. 19, 1997]

§ 520.88 Amoxicillin oral dosage forms.

§ 520.88a Amoxicillin trihydrate film-coated tablets.

(a) *Specifications*. Each tablet contains amoxicillin trihydrate equivalent to 50, 100, 150, 200, or 400 milligrams of amoxicillin.

(b) *Sponsor*. See No. 000069 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Dogs*—(i) *Amount*. 5 milligrams per pound of body weight, twice a day.

(ii) *Indications for use*. Treatment of infections of the respiratory tract (tonsillitis, tracheobronchitis), genitourinary tract (cystitis), gastrointestinal tract (bacterial gastroenteritis), and soft tissues (abscesses, lacerations, wounds), caused by susceptible strains of *Staphylococcus aureus*, *Streptococcus* spp., *Escherichia coli*, *Proteus mirabilis*, and bacterial dermatitis caused by *S. aureus*, *Streptococcus* spp., and *P. mirabilis*.

(iii) *Limitations*. Administer for 5 to 7 days or 48 hours after all symptoms have subsided. If no improvement is seen in 5 days, review diagnosis and change therapy. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Cats*—(i) *Amount*. 50 milligrams (5 to 10 milligrams per pound of body weight) once a day.

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(ii) *Indications for use*. Treatment of infections caused by susceptible organisms as follows: upper respiratory tract due to *S. aureus*, *Streptococcus* spp., and *E. coli*; genitourinary tract (cystitis) due to *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*; gastrointestinal tract due to *E. coli*; and skin and soft tissue (abscesses, lacerations, and wounds) due to *S. aureus*, *Streptococcus* spp., *E. coli*, and *Pasteurella multocida*.

(iii) *Limitations*. Administer for 5 to 7 days or 48 hours after all symptoms have subsided. If no improvement is seen in 5 days, review diagnosis and change therapy. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37319, Aug. 18, 1992, as amended at 60 FR 55658, Nov. 2, 1995]

§ 520.88b Amoxicillin trihydrate for oral suspension.

(a) *Specifications*. When reconstituted, each milliliter contains amoxicillin trihydrate equivalent to 50 milligrams of amoxicillin.

(b) *Sponsor*. See No. 000069 in § 510.600(c) of this chapter.

(1) *Conditions of use*—(i) *Dogs*—(A) *Amount*. 5 milligrams per pound of body weight twice daily.

(B) *Indications for use*. Treatment of infections caused by susceptible strains of organisms as follows: respiratory tract (tonsillitis, tracheobronchitis) caused by *Staphylococcus aureus*, *Streptococcus* spp., *Escherichia coli*, and *Proteus mirabilis*; genitourinary tract (cystitis) caused by *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*; gastrointestinal tract (bacterial gastroenteritis) caused by *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*; bacterial dermatitis caused by *S. aureus*, *Streptococcus* spp., and *P. mirabilis*; and soft tissues (abscesses, lacerations, and wounds) caused by *S. aureus*, *Streptococcus* spp., *E. coli*, and *P. mirabilis*.

(C) *Limitations*. Use for 5 to 7 days or 48 hours after all symptoms have subsided. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(ii) *Cats*—(A) *Amount*. 50 milligrams (5 to 10 milligrams per pound) once daily.

§ 520.88c

21 CFR Ch. I (4–1–03 Edition)

(B) *Indications for use.* Treatment of infections caused by susceptible strains of organisms as follows: upper respiratory tract due to *Staphylococcus* spp., *Streptococcus* spp., *Hemophilus* spp., *E. coli*, *Pasteurella* spp., and *P. mirabilis*; genitourinary tract (cystitis) due to *S. aureus*, *Streptococcus* spp., *E. coli*, *P. mirabilis*, and *Corynebacterium* spp.; gastrointestinal tract due to *E. coli*, *Proteus* spp., *Staphylococcus* spp., and *Streptococcus* spp.; skin and soft tissue (abscesses, lacerations, and wounds) due to *Staphylococcus* spp., *Streptococcus* spp., *E. coli*, and *Pasteurella multocida*.

(C) *Limitations.* Use for 5 to 7 days or 48 hours after all symptoms have subsided. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

(c) *Sponsor.* See Nos. 000093 and 000856 in § 510.600(c) of this chapter.

(1) *Conditions of use. Dogs*—(i) *Amount.* 5 milligrams per pound of body weight twice daily.

(ii) *Indications for use.* Treatment of bacterial dermatitis due to *S. aureus*, *Streptococcus* spp., *Staphylococcus* spp., and *E. coli*, and soft tissue infections (abscesses, wounds, lacerations) due to *S. aureus*, *Streptococcus* spp., *E. coli*, *P. mirabilis* and *Staphylococcus* spp.

(iii) *Limitations.* Use for 5 to 7 days. Continue for 48 hours after all symptoms have subsided. If no improvement is seen in 5 days, review diagnosis and change therapy. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[57 FR 37319, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992, as amended at 60 FR 55658, Nov. 2, 1995; 62 FR 13302, Mar. 20, 1997; 67 FR 67521, Nov. 6, 2002]

§ 520.88c Amoxicillin trihydrate oral suspension.

(a) *Specifications.* Each 0.8-milliliter dose contains amoxicillin trihydrate equivalent to 40 milligrams of amoxicillin.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.510 of this chapter.

(d) *Conditions of use. Swine*—(1) *Amount.* 40 milligrams orally, twice a day using a dosing pump.

(2) *Indications for use.* Treatment of baby pigs under 10 pounds for porcine colibacillosis caused by *Escherichia coli* susceptible to amoxicillin.

(3) *Limitations.* Treat animals for 48 hours after all symptoms have subsided but not beyond 5 days. Do not slaughter during treatment or for 15 days after latest treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37319, Aug. 18, 1992, as amended at 60 FR 55658, Nov. 2, 1995]

§ 520.88d Amoxicillin trihydrate soluble powder.

(a) *Specifications.* Each gram contains amoxicillin trihydrate equivalent to 115.4 milligrams of amoxicillin.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.38 of this chapter.

(d) *Conditions of use. Preruminating calves including veal calves*—(1) *Amount.* 400 milligrams per 100 pounds of body weight twice daily.

(2) *Indications for use.* Treatment of bacterial enteritis when due to susceptible *Escherichia coli* in preruminating calves including veal calves.

(3) *Limitations.* Administer by drench or by mixing in milk. Treatment should be continued for 48 hours after all symptoms have subsided but not to exceed 5 days. For use in preruminating calves including veal calves only, not for use in other animals which are raised for food production. Do not slaughter animals during treatment or for 20 days after the latest treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37319, Aug. 18, 1992; 57 FR 42623, Sept. 15, 1992; 58 FR 18304, Apr. 8, 1993, as amended at 60 FR 55658, Nov. 2, 1995; 62 FR 5525, Feb. 6, 1997]

§ 520.88e Amoxicillin trihydrate boluses.

(a) *Specifications.* Each bolus contains the equivalent of 400 milligrams of amoxicillin.

(b) *Sponsor.* See No. 000069 in § 510.600(c) of this chapter.