

(ii) *Indications for use.* For the control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), small strongyles (*Cyathostomum* spp., *Cylicocycclus* spp., *Cylicostephanus* spp., *Triodontophorus* spp.), pinworms (*Oxyuris equi*), and ascarids (*Parascaris equorum*) in horses.

(iii) *Limitations.* Administer orally by dose syringe or suitable plastic syringe. Do not use in horses intended for food. Consult a veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(2) *Cattle including dairy cows of breeding age*—(i) *Amount.* Administer orally 5 milligrams per kilogram of body weight (2.3 milligrams per pound).

(ii) *Indications for use.* For the removal and control of lungworm (*Dictyoacaulus viviparus*); stomach worm (adults)—brown stomach worm (*Ostertagia ostertagi*); stomach worms (adults and 4th-stage larvae)—barberpole worm (*Haemonchus contortus* and *H. placei*) and small stomach worm (*Trichostrongylus axei*); intestinal worms (adults and 4th-stage larvae)—hookworm (*Bunostomum phlebotomum*), threadnecked intestinal worm (*Nematodirus helvetianus*), small intestinal worm (*Cooperia punctata* and *C. oncophora*), bankrupt worm (*Trichostrongylus colubriformis*), and nodular worm (*Oesophagostomum radiatum*).

(iii) *Limitations.* Retreatment may be needed after 4 to 6 weeks. Cattle must not be slaughtered within 8 days following last treatment. Consult a veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(3) *Beef cattle*—(i) *Amount.* Administer orally 10 milligrams per kilogram of body weight.

(ii) For the removal and control of stomach worm (4th-stage inhibited larvae/type II ostertagiasis), *Ostertagia ostertagi*, and tapeworm, *Moniezia benedeni*.

(iii) *Limitations.* Retreatment may be needed after 4 to 6 weeks. Cattle must not be slaughtered within 8 days following last treatment. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(4) *Goats*—(i) *Amount.* Administer orally 5 milligrams per kilogram of body weight (2.3 milligrams per pound).

(ii) *Indications for use.* For the removal and control of stomach and intestinal worms *Haemonchus contortus* and *Ostertagia circumcincta*.

(iii) *Limitations.* Retreatment may be needed after 4 to 6 weeks. Goats must not be slaughtered for food within 6 days following last treatment. Do not use in lactating goats. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(e) *Special considerations.* Fenbendazole suspension 10 percent and approved forms of trichlorfon, when used concomitantly for treating the indications provided in paragraph (d) of this section and for treating infections of stomach bot as provided in § 520.2520, have been shown to be compatible and not to interfere with one another.

[42 FR 59069, Nov. 15, 1977; 43 FR 12311, Mar. 24, 1978. Redesignated at 44 FR 1375, Jan. 5, 1979, and amended at 46 FR 29464, June 2, 1981; 47 FR 15327, Apr. 9, 1982; 48 FR 42809, Sept. 20, 1983; 49 FR 1983, Jan. 17, 1984; 53 FR 40058, Oct. 13, 1988; 59 FR 26943, May 25, 1994; 61 FR 29478, June 11, 1996; 63 FR 63983, Nov. 18, 1998; 66 FR 47960, Sept. 17, 2001]

**§ 520.905b Fenbendazole granules.**

(a) *Specifications.* Each gram of granules contains 222 milligrams (mg) fenbendazole.

(b) *Sponsor.* See No. 057926 in § 510.600(c) of this chapter.

(c) *Special considerations.* See § 500.25 of this chapter.

(d) *Conditions of use*—(1) *Horses*—(i) *Amount.* 5 mg/kilogram (kg) for large strongyles, small strongyles, and pinworms; 10 mg/kg for ascarids.

(ii) *Indications for use.* For the control of infections of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), small strongyles, pinworms (*Oxyuris equi*), and ascarids (*Parascaris equorum*).

(iii) *Limitations.* Sprinkle the appropriate amount of drug on a small amount of the usual grain ration. Prepare for each horse individually. Withholding feed or water is not necessary. Retreat in 6 to 8 weeks if required. Do not use in horses intended for food.

(2) *Dogs*—(i) *Amount.* 50 mg/kg daily for 3 consecutive days.

(ii) *Indications for use.* For the removal of ascarids (*Toxocara canis*, *Toxascaris leonina*), hookworms (*Ancylostoma caninum*, *Uncinaria stenocephala*), whipworms (*Trichuris vulpis*), and tapeworms (*Taenia pisiformis*).

(iii) *Limitations.* Mix the appropriate amount of drug with a small amount of the usual food; dry dog food may require slight moistening to facilitate mixing. Medicated food must be fully consumed.

(3) *Zoo and wildlife animals—(i) Amount.* 10 mg/kg per day for 3 days.

(ii) *Indications for use.* For control of internal parasites of *Felidae* and *Ursidae* as follows:

(A) Lion (*Panthera leo*) and Tiger (*Panthera tigris*): Ascarid (*Toxocara cati*, *Toxascaris leonina*), Hookworm (*Ancylostoma* spp.).

(B) Cheetah (*Acinonyx jubatus*): Ascarid (*Toxocara cati*, *Toxascaris leonina*).

(C) Puma (*Felis concolor*), Panther (*Panthera* spp.), Leopard (*Panthera pardus*), Jaguar (*Panthera onca*): Ascarid (*Toxocara cati*, *Toxascaris leonina*), Hookworm (*Ancylostoma* spp.), Tapeworm (*Taenia hydatigena*, *T. krabbei*, *T. taeniaeformis*).

(D) Black Bear (*Ursus americanus*): Ascarid (*Baylisascaris transfuga*, *Toxascaris leonina*), Hookworm (*Ancylostoma caninum*), Tapeworm (*Taenia hydatigena*, *T. krabbei*).

(E) Polar Bear (*Ursus maritimus*) and Grizzly Bear (*Ursus horribilis*): Ascarid (*Baylisascaris transfuga*, *Toxascaris leonina*).

(iii) *Limitations.* Top dress or mix with a small portion of food. Must be fully consumed prior to feeding. Federal law restricts this drug to use by or on the order of a licensed veterinarian. Do not use 14 days before or during the hunting season.

[44 FR 1375, Jan. 5, 1979, as amended at 47 FR 15327, Apr. 9, 1982; 48 FR 50528, Nov. 2, 1983; 59 FR 35252, July 11, 1994; 66 FR 47960, Sept. 17, 2001; 67 FR 47450, July 19, 2002]

#### § 520.905c Fenbendazole paste.

(a) *Specifications.* The product is an aqueous paste containing 10 percent fenbendazole.

(b) *Sponsor.* See No. 057926 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.275 of this chapter.

(d) *Conditions of use—(1) Horses—(i)(a) Amount.* 2.3 milligrams per pound of body weight (one 2.5-gram fenbendazole syringe for a 1,100-pound horse). For foals and weanlings (less than 18 months of age), 4.6 milligrams per pound of body weight (one 2.5-gram fenbendazole syringe for each 550 pounds of body weight).

(b) *Indications for use.* For control of large strongyles (*Strongylus edentatus*, *S. equinus*, *S. vulgaris*), small strongyles, pinworms (*Oxyuris equi*), and ascarids (*Parascaris equorum*) in horses.

(c) *Limitations.* Retreatment at intervals of 6 to 8 weeks may be required. Do not use in horses intended for food. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

(ii)(a) *Amount.* 4.6 milligrams per pound of body weight (one 2.5-gram fenbendazole syringe for a 550-pound horse) daily for 5 days.

(b) *Indications for use.* For control of arteritis caused by the fourth stage larvae of *Strongylus vulgaris*.

(c) *Limitations.* Treatment should be initiated in the spring and repeated in 6 months. Do not use in horses intended for food. Consult your veterinarian for assistance in the diagnosis, treatment, and control of fourth stage larvae of *S. vulgaris*.

(iii)(a) *Amount.* 4.6 milligrams per pound of body weight (10 milligrams per kilogram) daily for 5 consecutive days.

(b) *Indications for use.* For treatment of encysted mucosal cyathostome (small strongyle) larvae including early third stage (hypobiotic), late third stage, and fourth stage larvae in horses.

(c) *Limitations.* (Consult your veterinarian for assistance in the diagnosis, treatment, and control of encysted mucosal cyathostomes). Do not use in horses intended for food.

(2) *Beef and dairy cattle—(i) Amount.* Administer orally 5 milligrams per kilogram of body weight (2.3 milligrams per pound).

(ii) *Indications for use.* For the removal and control of lungworm (*Dictyocaulus viviparus*), barberpole