

§ 522.1020

(A) *Indications for use.* For treatment of acute noninflammatory tissue edema.

(B) *Limitations.* Do not use in horses intended for food.

(3) *Cattle*—(i) *Amount.* 500 mg/animal once daily, intramuscularly or intravenously; or 250 mg/animal twice daily at 12-hour intervals, intramuscularly or intravenously.

(ii) *Indications for use.* For the treatment of physiological parturient edema of the mammary gland and associated structures.

(iii) *Limitations.* Treatment not to exceed 48 hours post-parturition. Milk taken during treatment and for 48 hours (four milkings) after the last treatment must not be used for food. Cattle must not be slaughtered for food within 48 hours following last treatment.

[66 FR 47961, Sept. 17, 2001, as amended at 67 FR 18086, Apr. 15, 2002]

§ 522.1020 Gelatin solution.

(a) *Specifications.* It is sterile and each 100 cubic centimeters contains 8 grams of gelatin in an 0.85 percent sodium chloride solution.

(b) *Sponsor.* See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used to restore circulatory volume and maintain blood pressure in animals being treated for shock.

(2) The exact dosage to be administered must be determined after evaluating the animal's condition and will vary according to the size of the animal and the degree of shock. A suggested dosage range for small animals such as dogs is 4 to 8 cubic centimeters per pound body weight. The suggested dosage range for large animals such as sheep, calves, cows, or horses is 2 to 4 cubic centimeters per pound of body weight. It is administered intravenously at a rate of 10 cubic centimeters per minute in small animals and 20 to 30 cubic centimeters per minute in large animals. The solution is administered aseptically and must be between 50 to 70 °F. when injected.

(3) A few animals will exhibit signs of allergic reaction. This solution can cause transient reversible nephrosis. This product is not intended to replace whole blood in cases of anemia and

21 CFR Ch. I (4–1–03 Edition)

should not be used in the presence of renal dysfunction. Unused portions remaining in bottles should be discarded.

(4) For use only by or on the order of a licensed veterinarian.

§ 522.1044 Gentamicin sulfate injection.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains gentamicin sulfate equivalent to either 5, 50, or 100 milligrams of gentamicin.

(b) *Sponsors.* (1) See No. 000061 in § 510.600(c) of this chapter for use of: 5-milligrams-per-milliliter solution in swine as in paragraph (d)(4) of this section, 50-milligrams-per-milliliter solution in dogs and cats as in paragraph (d)(1) of this section, 50- and 100-milligrams-per-milliliter solution in chickens and turkeys as in paragraphs (d)(2) and (d)(3) of this section.

(2) [Reserved]

(3) See No. 000010 for use of 50 milligrams-per-milliliter solution in dogs as in paragraph (d)(5) of this section.

(4) See No. 059130 for use of 100 milligram-per-milliliter solution in turkeys as in paragraph (d)(2) of this section and in chickens as in paragraph (d)(3) of this section.

(c) *Related tolerances.* See § 556.300 of this chapter.

(d) *Conditions of use*—(1) *Dogs and cats*—(i) *Amount.* Two milligrams of gentamicin per pound of body weight, twice daily on the first day, once daily thereafter, using a 50 milligram-per-milliliter solution.

(ii) *Indications for use*—(a) *Dogs.* For the treatment of infections of urinary tract (cystitis, nephritis), respiratory tract (tonsillitis, pneumonia, tracheobronchitis), skin and soft tissue (pyodermatitis, wounds, lacerations, peritonitis).

(b) *Cats.* For the treatment of infections of urinary tract (cystitis, nephritis), respiratory tract (pneumonitis, pneumonia, upper respiratory tract infections), skin and soft tissue (wounds, lacerations, peritonitis), and as supportive therapy for secondary bacterial infections associated with panleucopenia.

(iii) *Limitations.* Administer intramuscularly or subcutaneously. If response is not noted after 7 days, the antibiotic sensitivity of the infecting

organism should be retested. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Turkeys*—(i) *Amount*. One milligram of gentamicin per 0.2 milliliter dose, using the 50- or 100-milligrams-per-milliliter product diluted with sterile saline to a concentration of 5 milligrams-per-milliliter.

(ii) *Indications for use*. As an aid in the prevention of early mortality due to Arizona paracolon infections susceptible to gentamicin.

(iii) *Limitations*. For 1- to 3-day old turkey poults. Administer subcutaneously in the neck. Injected poults must not be slaughtered for food for at least 9 weeks after treatment.

(3) *Chickens*—(i) *Amount*. 0.2 milligram of gentamicin per 0.2 milliliter dose, using the 50- or 100-milligrams-per-milliliter product diluted with sterile saline to a concentration of 1.0 milligram-per-milliliter.

(ii) *Indications for use*. In day-old chickens, for prevention of early mortality caused by *Escherichia coli*, *Salmonella typhimurium*, and *Pseudomonas aeruginosa* that are susceptible to gentamicin.

(iii) *Limitations*. For use in day-old chickens only. Administer aseptically, injecting the diluted product subcutaneously in the neck. Do not slaughter treated animals for food for at least 5 weeks after treatment.

(4) *Swine*—(i) *Amount*. 5 milligrams of gentamicin as a single intramuscular dose using 5 milligram-per-milliliter solution.

(ii) *Indications for use*. In piglets up to 3 days old for treatment of porcine colibacillosis caused by strains of *E. coli* sensitive to gentamicin.

(iii) *Limitations*. For single intramuscular dose in pigs up to 3 days of age only. Do not slaughter treated animals for food for at least 40 days following treatment.

(5) *Dogs*—(i) *Amount*. 2 milligrams of gentamicin per pound of body weight, twice daily on the first day, then once daily.

(ii) *Indications for use*. For use in the treatment of urinary tract infections (cystitis) caused by *Proteus mirabilis*, *Escherichia coli*, and *Staphylococcus aureus*.

(iii) *Limitations*. Administer intramuscularly or subcutaneously. If no improvement is seen after 3 days, treatment should be discontinued and the diagnosis reevaluated. Treatment not to exceed 7 days. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[43 FR 1942, Jan. 13, 1978, as amended at 48 FR 791, Jan. 7, 1983; 51 FR 15606, Apr. 25, 1986; 52 FR 7832, Mar. 13, 1987; 53 FR 40727, Oct. 18, 1988; 60 FR 29985, June 7, 1995; 61 FR 24441, May 15, 1996; 62 FR 45157, Aug. 26, 1997; 63 FR 59714, Nov. 5, 1998; 63 FR 68182, Dec. 10, 1998; 65 FR 45877, July 26, 2000]

§ 522.1055 Gleptoferron injection.

(a) *Specifications*. Each milliliter contains the equivalent of 200 milligrams of elemental iron as gleptoferron (complex of ferric hydroxide and dextran glucoheptonic acid), and 0.5 percent phenol as a preservative.

(b) *Sponsor*. See 062408 in § 510.600(c) of this chapter.

(c) *Conditions of use*. It is used in baby pigs as follows:

(1) For prevention of iron deficiency anemia, administer 200 milligrams of elemental iron intramuscularly on or before 3 days of age.

(2) For treatment of iron deficiency anemia, administer 200 milligrams of elemental iron intramuscularly.

[45 FR 61288, Sept. 16, 1980, as amended at 61 FR 18672, Apr. 29, 1996]

§ 522.1066 Glycopyrrolate injection.

(a) *Specifications*. Each milliliter of aqueous solution contains 0.2 milligram of glycopyrrolate.

(b) *Sponsor*. See No. 000856 in § 510.600(c) of this chapter.

(c) *Conditions of use*. (1) It is indicated as a preanesthetic agent in dogs and cats.

(2) It is administered intravenously, intramuscularly, or subcutaneously in dogs and intramuscularly in cats at a dosage level of 5 micrograms per pound of body weight (0.25 milliliter per 10 pounds of body weight).

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[48 FR 21567, May 13, 1983, as amended at 67 FR 67521, Nov. 6, 2002]