

§ 522.1260

21 CFR Ch. I (4-1-03 Edition)

(c) *Conditions of use*—(1) *Amount*. 2 milliliters per 100 pounds of body weight, subcutaneously in the neck.

(2) *Indications for use*. (i) The 13.65 percent injection is used as an anthelmintic in cattle for treatment of the following parasites: stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*, *Chabertia*), and lungworms (*Dictyocaulus*).

(ii) The 18.2 percent injection is used as an anthelmintic in cattle for treatment of the following parasites: stomach worms (*Haemonchus*, *Trichostrongylus*, *Ostertagia*), intestinal worms (*Trichostrongylus*, *Cooperia*, *Nematodirus*, *Bunostomum*, *Oesophagostomum*) and lungworms (*Dictyocaulus*).

(3) *Limitations*. Do not administer more than 10 milliliters per site. Cattle that are severely parasitized or maintained under conditions of constant helminth exposure may require retreatment within 2 to 4 weeks after first treatment. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. Consult your veterinarian before using in severely debilitated animals or animals under severe stress. Do not administer to cattle within 7 days of slaughter. Do not administer to dairy animals of breeding age.

[43 FR 20489, May 12, 1978, as amended at 43 FR 29289, July 7, 1978; 43 FR 60895, Dec. 29, 1978; 47 FR 10807, Mar. 12, 1982; 62 FR 61625, Nov. 19, 1997; 65 FR 61090, Oct. 16, 2000; 67 FR 63055, Oct. 10, 2002]

§ 522.1260 **Lincomycin.**

(a) *Specifications*. Each milliliter of solution contains lincomycin hydrochloride monohydrate equivalent to 25, 50, 100, or 300 milligrams (mg) of lincomycin.

(b) *Sponsors*. See sponsors in § 510.600(c) of this chapter for uses as in paragraph (e) of this section.

(1) No. 000009 for uses as in paragraph (e) of this section.

(2) No. 046573 for use as in paragraph (e)(2) of this section.

(c) *Special considerations*. When common labeling for use of the drug in dogs, cats, and swine is included with

the drug, all such uses are subject to the labeling requirements of § 201.105 of this chapter.

(d) *Related tolerances*. See § 556.360 of this chapter.

(e) *Conditions of use*. It is used for animals as follows:

(1) *Dogs and cats*—(i) *Amount*. 5 mg per pound (lb) of body weight twice daily or 10 mg/lb body weight once daily by intramuscular injection; 5 to 10 mg/lb body weight one or two times daily by slow intravenous injection.

(ii) *Indications for use*. Infections caused by Gram-positive organisms, particularly streptococci and staphylococci.

(iii) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) *Swine*—(i) *Amount*. 5 mg/lb body weight once daily by intramuscular injection for 3 to 7 days.

(ii) *Indications for use*. Treatment of infectious arthritis and mycoplasma pneumonia.

(iii) *Limitations*. Do not treat within 48 hours of slaughter.

[40 FR 13858, Mar. 27, 1975, as amended at 50 FR 31351, Aug. 2, 1985; 67 FR 34388, May 14, 2002]

§ 522.1289 **Lufenuron suspension.**

(a) *Specifications*. Each milliliter of sterile aqueous suspension contains 10 milligrams of lufenuron.

(b) *Sponsor*. See No. 058198 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(1) *Cats*—(i) *Amount*. 10 milligrams per kilogram (4.5 milligrams per pound) of body weight every 6 months, subcutaneously.

(ii) *Indications for use*. For use in cats 6 weeks of age and older, for control of flea populations. Lufenuron controls flea populations by preventing the development of flea eggs and does not kill adult fleas. Concurrent use of insecticides may be necessary for adequate control of adult fleas.

(iii) *Limitations*. For subcutaneous use in cats only. The safety of this product in reproducing animals has not been established. Do not use in dogs. Federal law restricts this drug to use by or on the order of a licensed veterinarian.