

(2) *Indications for use.* For synchronization of estrus/ovulation in cycling beef cattle and non-lactating dairy heifers.

(3) *Limitations.* Insert implant subcutaneously in the ear only; then immediately inject solution intramuscularly only. Counting the day of implantation as day 1, remove the implant on day 10. Collect all implants as they are removed and burn them. While animals are restrained for artificial insemination, avoid other treatments such as vaccinations, dipping, pour-on grub and louse prevention, spraying, etc. When inseminating without estrus detection, the entire treated group should be started at 48 hours after the last implant has been removed and should be completed within 6 hours. Where estrus detection is preferred, insemination should be approximately 12 hours after first detection of estrus. Those that do not conceive can be re-bred when they return to estrus approximately 17 to 25 days after implant removal. Do not use in cows producing milk for human consumption.

[47 FR 55477, Dec. 10, 1982, as amended at 48 FR 49656, Oct. 27, 1983; 51 FR 33592, Sept. 22, 1986; 54 FR 1165, Jan. 12, 1989]

§ 522.863 Ethylisobutrazine hydrochloride injection.

(a) *Specifications.* The drug is a sterile aqueous solution. Each milliliter contains 50 milligrams of ethylisobutrazine hydrochloride.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) It is used in dogs as a tranquilizer.¹

(2) It is administered intramuscularly at a dosage level of 2 to 5 milligrams of ethylisobutrazine hydrochloride per pound of body weight for profound tranquilization. It is administered intravenously at a dosage level of 1 to 2 milligrams of ethylisobutrazine hydrochloride per pound of body weight to effect.¹

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(3) It is not to be used in conjunction with organophosphates and/or procaine hydrochloride because phenothiazines may potentiate the toxicity of organophosphates and the activity of procaine hydrochloride.¹

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13858, Mar. 27, 1975, as amended at 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§ 522.883 Etorphine hydrochloride injection.

(a) *Chemical name.* 6,7,8,14 - tetrahydro - alpha - methyl - alpha - propyl - 6,14 - endo-ethenooripavine-alpha-methanol hydrochloride.

(b) *Specifications.* Each milliliter of etorphine hydrochloride injection, veterinary, contains 1 mg of etorphine hydrochloride in sterile aqueous solution.

(c) *Sponsors.* See No. 053923 in § 510.600(c) of this chapter.

(d) *Conditions of use.* (1) The drug is used for the immobilization of wild and exotic animals.

(2) It is administered intramuscularly by hand syringe or syringe dart at a suitable dosage level depending upon the species.

(3) Do not use the drug unless diprenorphine hydrochloride injection, veterinary, as provided for in § 522.723, is available for use in reversing the effects of etorphine hydrochloride injection, veterinary.

(4) For use in wild or exotic animals only. Do not use in domestic food-producing animals. Do not use 30 days before, or during, the hunting season in free-ranging wild animals that might be used for food.

(5) Federal law restricts this drug to use by or on the order of a licensed veterinarian. Distribution is restricted to veterinarians engaged in zoo and exotic animal practice, wildlife management programs, and researchers.

[40 FR 13858, Mar. 27, 1975, as amended at 48 FR 16241, Apr. 15, 1983; 61 FR 260, Jan. 4, 1996]

§ 522.900 Euthanasia solution.

(a) [Reserved]

(b)(1) *Specifications.* Each milliliter of nonsterile solution contains 390 milligrams of pentobarbital sodium and 50 milligrams of phenytoin sodium.