

## § 556.1

556.530	Prednisone.
556.540	Progesterone.
556.550	Propylparaben.
556.560	Pyrantel tartrate.
556.570	Ractopamine.
556.580	Robenidine hydrochloride.
556.590	Salicylic acid.
556.592	Salinomycin.
556.597	Semduramicin.
556.600	Spectinomycin.
556.610	Streptomycin.
556.620	Sulfabromomethazine sodium.
556.625	Sodium sulfachloropyrazine monohydrate.
556.630	Sulfachlorpyridazine.
556.640	Sulfadimethoxine.
556.650	Sulfaethoxyypyridazine.
556.660	Sulfamerazine.
556.670	Sulfamethazine.
556.680	Sulfanitran.
556.685	Sulfaquinoxaline.
556.690	Sulfathiazole.
556.700	Sulfomyxin.
556.710	Testosterone propionate.
556.720	Tetracycline.
556.730	Thiabendazole.
556.735	Tilmicosin.
556.738	Tiamulin.
556.739	Trenbolone.
556.740	Tylosin.
556.741	Tripelennamine.
556.750	Virginiamycin.
556.760	Zeranol.
556.770	Zoalene.

AUTHORITY: 21 U.S.C. 342, 360b, 371.

SOURCE: 40 FR 13942, Mar. 27, 1975, unless otherwise noted.

### Subpart A—General Provisions

#### § 556.1 General considerations; tolerances for residues of new animal drugs in food.

(a) Tolerances established in this part are based upon residues of drugs in edible products of food-producing animals treated with such drugs. Consideration of an appropriate tolerance for a drug shall result in a conclusion either that:

(1) Finite residues will be present in the edible products—in which case a finite tolerance is required; or

(2) It is not possible to determine whether finite residues will be incurred but there is reasonable expectation that they may be present—in which case a tolerance for negligible residue is required; or

(3) The drug induces cancer when ingested by man or animal or, after tests which are appropriate for the evaluation of the safety of such drug, has

## 21 CFR Ch. I (4–1–03 Edition)

been shown to induce cancer in man or animal; however, such drug will not adversely affect the animals for which it is intended, and no residue of such drug will be found by prescribed methods of analysis in any edible portion of such animals after slaughter or in any food yielded by or derived from the living animal—in which case the accepted method of analysis shall be published or cited, if previously published and available elsewhere, in this part; or

(4) It may or may not be possible to determine whether finite residues will be incurred but there is no reasonable expectation that they may be present—in which case the establishment of a tolerance is not required; or

(5) The drug is such that it may be metabolized and/or assimilated in such form that any possible residue would be indistinguishable from normal tissue constituents—in which case the establishment of a tolerance is not required.

(b) No tolerance established pursuant to paragraph (a)(1) of this section will be set at any level higher than that reflected by the permitted use of the drug.

(c) Any tolerance required pursuant to this section will, in addition to the toxicological considerations, be conditioned on the availability of a practicable analytical method to determine the quantity of residue. Such method must be sensitive to and reliable at the established tolerance level or, in certain instances, may be sensitive at a higher level where such level is also deemed satisfactory and safe in light of the toxicity of the drug residue and of the unlikelihood of such residue's exceeding the tolerance.

### Subpart B—Specific Tolerances for Residues of New Animal Drugs

#### § 556.20 2-Acetylamino-5-nitrothiazole.

A tolerance of 0.1 part per million is established for negligible residues of 2-acetylamino-5-nitrothiazole in the edible tissues of turkeys.

#### § 556.30 Aklomide.

Tolerances are established for combined residues of aklomide (2-chloro-4-nitrobenzamide) and its metabolite (4-

amino-2-chlorobenzamide) in uncooked edible tissues of chickens as follows:

(a) 4.5 parts per million in liver and muscle.

(b) 3 parts per million in skin with fat.

**§ 556.34 Albendazole.**

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of albendazole is 5 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Cattle*. A tolerance is established for albendazole 2-aminosulfone (marker residue) in liver (target tissue) of 0.2 part per million and in muscle of 0.05 part per million.

(2) *Sheep*. A tolerance is established for albendazole 2-aminosulfone (marker residue) in liver (target tissue) of 0.25 part per million and in muscle of 0.05 part per million.

[64 FR 1504, Jan. 11, 1999]

**§ 556.38 Amoxicillin.**

A tolerance of 0.01 part per million is established for negligible residues of amoxicillin in milk and in the uncooked edible tissues of cattle.

[49 FR 45422, Nov. 16, 1984]

**§ 556.40 Ampicillin.**

A tolerance of 0.01 p/m is established for negligible residues of ampicillin in the uncooked edible tissues of swine and cattle and in milk.

**§ 556.50 Amprolium.**

Tolerances are established as follows for residues of amprolium (1-(4-amino-2-*n*-propyl-5-pyrimidinylmethyl)-2-picolinium chloride hydrochloride):

(a) In the edible tissues and in eggs of chickens and turkeys:

(1) 1 part per million in uncooked liver and kidney.

(2) 0.5 part per million in uncooked muscle tissue.

(3) In eggs:

(i) 8 parts per million in egg yolks.

(ii) 4 parts per million in whole eggs.

(b) In the edible tissues of calves:

(1) 2.0 parts per million in uncooked fat.

(2) 0.5 part per million in uncooked muscle tissue, liver, and kidney.

(c) In the edible tissues of pheasants:

(1) 1 part per million in uncooked liver.

(2) 0.5 part per million in uncooked muscle.

[40 FR 13942, Mar. 27, 1975, as amended at 50 FR 18472, May 1, 1985]

**§ 556.52 Apramycin.**

A tolerance of 0.1 part per million is established for parent apramycin (marker residue) in kidney (target tissue) of swine. The acceptable daily intake (ADI) for total residues of apramycin is 25 micrograms per kilogram of body weight per day.

[62 FR 40933, July 31, 1997]

**§ 556.60 Arsenic.**

Tolerances for total residues of combined arsenic (calculated as As) in food are established as follows:

(a) In edible tissues and in eggs of chickens and turkeys:

(1) 0.5 part per million in uncooked muscle tissue.

(2) 2 parts per million in uncooked edible by-products.

(3) 0.5 part per million in eggs.

(b) In edible tissues of swine:

(1) 2 parts per million in uncooked liver and kidney.

(2) 0.5 part per million in uncooked muscle tissue and by-products other than liver and kidney.

**§ 556.70 Bacitracin.**

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of bacitracin is 0.05 milligram per kilogram of body weight per day.

(b) *Tolerances*. The tolerance for residues of bacitracin from zinc bacitracin or bacitracin methylene disalicylate in uncooked edible tissues of cattle, swine, chickens, turkeys, pheasants, and quail, and in milk and eggs is 0.5 part per million.

[65 FR 70791, Nov. 28, 2000]

**§ 556.90 Buquinolate.**

Tolerances are established for residues of buquinolate as follows:

(a) In edible tissues of chickens:

(1) 0.4 part per million in uncooked liver, kidney, and skin with fat.

(2) 0.1 part per million in uncooked muscle.

(b) In eggs: