

**Food and Drug Administration, HHS**

**§ 558.128**

feeds containing less than 15 percent crude protein.

[40 FR 13959, Mar. 27, 1975, as amended at 40 FR 45164, Oct. 1, 1975; 40 FR 57798, Dec. 12, 1975; 42 FR 761, Jan. 4, 1977; 51 FR 7396, Mar. 3, 1986; 63 FR 59216, Nov. 3, 1998; 66 FR 47963, Sept. 17, 2001]

**§ 558.120 Carbarsonone (not U.S.P.).**

(a) *Approvals.* Type A medicated articles: (1) 37.5 percent to 046573 in § 510.600(c) of this chapter.

(2) 25 percent carbarsonone and 5 grams per pound bacitracin (as bacitracin methylene disalicylate) to 046573 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.60 of this chapter.

(c) [Reserved]

(d) *Conditions of use.* (1) It is used for turkeys as follows:

(i) *Grams per ton.* 227 to 340.5 (0.025 to 0.0375 percent).

(a) *Indications for use.* As an aid in the prevention of blackhead.

(b) *Limitations.* Feed continuously beginning 2 weeks before blackhead is expected and continue as long as prevention is needed; withdraw 5 days before slaughter; as sole source of organic arsenic.

(ii) *Grams per ton.* 227 to 340.5 (0.025 to 0.0375 percent) carbarsonone plus 10 grams per ton bacitracin from bacitracin methylene disalicylate.

(a) *Indications for use.* As an aid in the prevention of blackhead; for increased rate of weight gain.

(b) *Limitations.* Feed continuously beginning 2 weeks before blackhead is expected and continue as long as prevention is needed; withdraw 5 days before slaughter; as sole source of organic arsenic.

(iii) *Grams per ton.* 227 to 340.5 (0.025 to 0.0375 percent) carbarsonone plus 4 to 45 grams per ton bacitracin from bacitracin zinc.

(a) *Indications for use.* As an aid in the prevention of blackhead, increased rate of weight gain, and improved feed efficiency.

(b) *Limitations.* Feed continuously as sole ration. Withdraw 5 days before slaughter. As sole source of organic arsenic; as bacitracin zinc provided by No. 046573 in § 510.600(c) of this chapter.

(2) Carbarsonone (not U.S.P.) may be used in accordance with the provisions of this section in the combinations provided as follows:

(i) Zoalene in accordance with § 558.680.

(ii) Amprolium as in § 558.55.

(iii) Bambermycins as in § 558.95.

[41 FR 10995, Mar. 15, 1976, as amended at 42 FR 18617, Apr. 8, 1977; 46 FR 46797, Sept. 22, 1981; 48 FR 2758, Jan. 21, 1983; 51 FR 7396, Mar. 3, 1986; 52 FR 2687, Jan. 26, 1987; 53 FR 20843, June 7, 1988; 57 FR 7652, Mar. 4, 1992; 61 FR 515, Jan. 8, 1996; 61 FR 18082, Apr. 24, 1996; 62 FR 61011, Nov. 14, 1997; 63 FR 27845, May 21, 1998; 66 FR 46706, Sept. 7, 2001]

**§ 558.128 Chlortetracycline.**

(a) *Specifications.* Type A medicated articles containing either chlortetracycline calcium complex equivalent to chlortetracycline hydrochloride or, for products intended for use in milk replacer, chlortetracycline hydrochloride.

(b) *Approvals.* See sponsors in § 510.600(c) of this chapter for use as in paragraph (e) of this section.

(1) Nos. 046573, 053389, and 066104: 50 to 100 grams per pound (g/lb) of Type A medicated article.

(2) No. 017519: 50 g/lb of Type A medicated article.

(c) *Related tolerances.* See § 556.150 of this chapter.

(d) *Special considerations.* (1) In milk replacers or starter feed; include on labeling the warning: "A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal."

(2) Manufacture for use in free-choice feeds as in paragraph (e)(4)(iii) of this section must conform to § 510.455 of this chapter.

(3) When manufactured for use as in paragraph (e)(5)(iv) of this section, include on labeling the warning: "Psittacosis, avian chlamydiosis, or ornithosis is a reportable communicable disease, transmissible between wild and domestic birds, other animals, and man. Contact appropriate public health and regulatory officials."

(e) *Conditions of use*—(1) *Chickens.* It is used as follows: